arXiv:hep-ph/9808421v1 26 Aug 1998

LU TP 98-17
UWThPh-1998-48
hep-ph/9808421
August 1998

Double Chiral Logs*

J. Bijnens[1], G. Colangelo[2 #] and G. Ecker[3]

[1] Dept. of Theor. Phys., Univ. Lund, Sölvegatan 14A, S–22362 Lund, Sweden
[2] INFN – Laboratori Nazionali di Frascati, P.O. Box 13, I–00044 Frascati, Italy
[3] Inst. Theor. Phys., Univ. Wien, Boltzmanng. 5, A–1090 Wien, Austria

Abstract

We determine the full structure of the leading (double-pole) divergences of $O(p^6)$ in the meson sector of chiral perturbation theory. The field theoretic basis for this calculation is described. We then use an extension of this result to determine the p^6 contributions containing double chiral logarithms (L^2), single logarithms times p^4 constants ($L \times L_i^r$) and products of two p^4 constants ($L_i^r \times L_j^r$) for F_π, F_K/F_π, K_{l3} and K_{e4} form factors. Numerical results are presented for these quantities.

* Work supported in part by TMR, EC-Contract No. ERBFMRX-CT980169 (EURODAΦNE).

[#] Address after september 1 1998: Institut für Theoretische Physik der Universität Zürich, Winterthurerstr. 190, CH–8057 Zürich–Irchel.

1. Chiral logs arise in the process of renormalization. For instance, in dimensional regularization a renormalization scale μ is introduced to ensure the correct dimension of amplitudes in d dimensions. One-loop divergences then always appear in the scale independent combination

$$\mathcal{X} = \frac{\mu^{d-4}}{(4\pi)^2}\left[\frac{1}{d-4} + \frac{1}{2}\ln\frac{M^2}{\mu^2} + \ldots\right] \tag{1}$$

with a characteristic meson mass M. Renormalization consists in canceling the pole in $d-4$ by the divergent part of the tree-level amplitude of $O(p^4)$ and replacing it by a combination of scale dependent low-energy constants from $\mathcal{L}_4$, the next-to-leading term in the low-energy expansion of the effective chiral Lagrangian $\mathcal{L}_{\mathrm{eff}} = \mathcal{L}_2 + \mathcal{L}_4 + \mathcal{L}_6 + \ldots$. In many cases, the chiral logs $\ln M^2/\mu^2$ make sizable contributions for a typical scale μ of $O(M_\rho)$.

At $O(p^6)$, the leading divergences are double poles accompanied by double chiral logs, the leading infrared singularities of $O(p^6)$. The double logs are again numerically important in general, e.g., for S-wave threshold parameters in $\pi\pi$ scattering [1, 2]. As a by-product of the complete renormalization of the generating functional of $O(p^6)$ [3], we present here the double chiral logs in full generality for chiral $SU(n)$. As will be shown below, the double logs (L^2) come together with terms of the form $L \times L_i^r$ and products $L_i^r \times L_j^r$ where the L_i^r are the renormalized low-energy constants of $O(p^4)$ (we use the $SU(3)$ notation here for simplicity). It is then very natural to include such terms in the numerical analysis especially because they are often comparable to or even bigger than the proper double-log terms.

The purpose of this letter is twofold. First, our general double-pole divergence Lagrangian for $SU(n)$ may serve as a check of existing and yet to be performed two-loop calculations of $O(p^6)$. Secondly, the numerical analysis may provide hints as to where large p^6 corrections are to be expected. Of course, the partial p^6 results presented here are not a substitute for the full expressions of $O(p^6)$. But once the double-pole divergence Lagrangian has been determined from a one-loop calculation (see below), the numerical applications come at almost no cost compared to the full p^6 calculations.

2. In a mass independent regularization scheme like dimensional regularization, the divergent parts are polynomials in masses and external momenta. In the present case, the coefficients of those polynomials are renormalized by the general chiral Lagrangian of $O(p^6)$. Therefore, the divergences themselves can be cast into the form of $\mathcal{L}_6$ with divergent coefficients that receive contributions from the diagrams shown in Fig. 1.

There are two types of diagrams in Fig. 1, reducible and irreducible ones, both divergent in general. However, as will be shown below, one can always choose $\mathcal{L}_4$ in such a way as to make the sum of the reducible diagrams c,e,g finite. Even so, these diagrams give rise to double logs in general. These double logs are completely determined by the tree-level contribution g proportional to products of low-energy constants of $O(p^4)$.

The tree-level diagram f being trivial, we are left with the irreducible loop diagrams a,b,d in Fig. 1. Starting with the two-loop diagrams a, b, the corresponding



Figure 1: Diagrams contributing to the generating functional of $O(p^6)$. The propagators and vertices carry the full tree structure associated with the lowest–order Lagrangian $\mathcal{L}_2$. Normal vertices are from $\mathcal{L}_2$, crossed circles denote vertices from $\mathcal{L}_4$ and the square in diagram f stands for a vertex from $\mathcal{L}_6$.

divergences in a given coefficient of $\mathcal{L}_6$ are of the general form

$$\begin{aligned} & x(d)\frac{\mu^{2(d-4)}}{(4\pi)^4}\left[\frac{1}{d-4}+\frac{1}{2}\ln\frac{M^2}{\mu^2}+\ldots\right]^2 \\ = \quad & x(d)\frac{\mu^{2(d-4)}}{(4\pi)^4}\left[\frac{1}{(d-4)^2}+\frac{\ln M^2/\mu^2}{d-4}+\frac{1}{4}(\ln\frac{M^2}{\mu^2})^2+\ldots\right] \end{aligned} \tag{2}$$

where $x(d)$ is a function of d that depends on the specific coefficient under consideration.

The problematic piece in (2) is the nonlocal divergence $x_0 \ln M^2/\mu^2/((4\pi)^4(d-4))$ that cannot be renormalized with a local action of $O(p^6)$. Here, x_0 is the leading coefficient in the Taylor expansion of $x(d)$ around $d=4$: $x(d)=x_0+x_1(d-4)+\cdots$. Therefore, in accordance with general theorems of renormalization theory [4], the nonlocal divergence must cancel with other contributions. Since the sum of reducible diagrams in Fig. 1 is finite with our choice of $\mathcal{L}_4$, the cancellation can only come from diagram d that is proportional to the low-energy constants L_i . These constants are themselves divergent (to renormalize the one-loop divergences). In d dimensions, they can be written

$$L_i(d)=\mu^{d-4}\left[\frac{\Gamma_i}{(4\pi)^2(d-4)}+L_i^r(\mu)+\ldots\right] \tag{3}$$

with known coefficients Γ_i and renormalized constants $L_i^r(\mu)$ [5]. The divergences due to the one-loop diagram d in Fig. 1 are then given by the product of (1) and

(3) with appropriate coefficients $y_i(d)$:

$$
\begin{aligned}
-\frac{y_i(d)}{2}\mathcal{X}L_i(d) &= -\frac{y_i(d)}{2}\frac{\mu^{2(d-4)}}{(4\pi)^4}\left[\frac{\Gamma_i}{(d-4)^2}+\frac{\Gamma_i \ln M^2/\mu^2}{2(d-4)}+\frac{(4\pi)^2 L_i^r(\mu)}{d-4}\right.\\
&+ \left.\frac{(4\pi)^2}{2}L_i^r(\mu)\ln M^2/\mu^2+\ldots\right] . \qquad (4)
\end{aligned}
$$

Summation over i is implied and we have chosen the numerical factor $-1/2$ in (4) to conform to the notation of Ref. [2]. Cancellation of the nonlocal divergences in (2) and (4) is guaranteed by Weinberg's conditions [6]

$$
x_0 = \frac{1}{4} y_{i0}\Gamma_i \; , \qquad (5)
$$

with y_{i0} the leading coefficient in the Taylor expansion of $y_i(d) = y_{i0} + (d-4)y_{i1} + \ldots$. In the chiral Lagrangian of $O(p^6)$ for $SU(n)$, there are 115 independent terms (including three contact terms) [3]. We have verified the corresponding 115 relations (5) by explicit calculation of both two- and one-loop divergences.

The divergences in the sum of (2) and (4) now have the proper local form to be renormalized by the counterterm action of $O(p^6)$ represented by diagram f in Fig. 1. After renormalization, we are left with finite parts

$$
\frac{x_0}{4(4\pi)^4}(\ln\frac{M^2}{\mu^2})^2 - \frac{y_{i0}L_i^r(\mu)}{4(4\pi)^2}\ln\frac{M^2}{\mu^2}+\ldots \qquad (6)
$$

The terms listed explicitly are the double-log contribution and all products of renormalized low-energy constants with a chiral log. With the help of (5), these two terms can be combined to give

$$
\frac{y_{i0}}{16}\left[\Gamma_i L^2 - 4L_i^r(\mu)L\right] := -\frac{y_{i0}}{16}k_i \qquad , \qquad L = \frac{1}{(4\pi)^2}\ln\frac{M^2}{\mu^2} \; . \qquad (7)
$$

The combinations k_i [1, 2] always appear together in the renormalized coefficients. They are process independent generalizations of the double logs and their coefficients are completely calculable from the one-loop diagram d in Fig. 1. The main reason for using the k_i together with the products $L_i^r \times L_j^r$ in numerical applications instead of only the double logs is a practical one. Very often, in particular for chiral $SU(3)$ for the conventional choices of M/μ, the additional terms are numerically at least as important as the double logs themselves.

Before discussing the actual calculation in more detail, we present the final result below. Since up to 11 of the k_i can appear in each of the 115 coefficients of $\mathcal{L}_6$, the complete Lagrangian as a function of the k_i cannot be reproduced here. Instead, we only write down the proper double-log Lagrangian for chiral $SU(n)$ in Minkowski space, i.e. dropping the terms[1] of the form $L_i^r(\mu)L$. We also drop all terms that do not contribute to processes with up to four mesons or one external field and three mesons.

---

[1] The full expression with all k_i and all terms can be obtained from the authors on request.

$$\begin{aligned}
\mathcal{L}_{\rm double-log} = \frac{L^2}{F^2}\{ & -(7/64+5/1152\, n^2)\langle A h_{\mu\nu}h^{\mu\nu}\rangle - 41/2304\, n\, \langle A\rangle\langle h_{\mu\nu}h^{\mu\nu}\rangle \\
& +3/64\,\langle h_{\mu\nu}u_\rho h^{\mu\nu}u^\rho\rangle + 1/288\,n\,\langle h_{\mu\nu}u_\rho\rangle\langle h^{\mu\nu}u^\rho\rangle - 1/16\,\langle h_{\mu\nu}\,(u_\rho h^{\mu\rho}u^\nu + u^\nu h^{\mu\rho}u_\rho)\rangle \\
& -1/384\,n\,\langle h_{\mu\nu}u_\rho\rangle\langle h^{\mu\rho}u^\nu\rangle + 1/144\,n^2\,\langle A^2\chi_+\rangle + 1/18\,n\,\langle A^2\rangle\langle\chi_+\rangle \\
& -1/64\,n\,\langle A\rangle\langle A\chi_+\rangle - 1/128\,\langle A\rangle^2\langle\chi_+\rangle - (3/32 - 17/1152\,n^2)\langle A u_\mu\chi_+u^\mu\rangle \\
& +(3/32-1/72\,n^2)\langle\chi_+u_\mu u_\nu u^\mu u^\nu\rangle - 1/18\,n\,\langle\chi_+\rangle\langle u_\mu u_\nu u^\mu u^\nu\rangle \\
& +1/32\,n\,\langle\chi_+u_\mu u_\nu\rangle\langle u^\mu u^\nu\rangle + 1/64\,\langle\chi_+\rangle\langle u_\mu u_\nu\rangle^2 - 19/2304\,n^2\,\langle\chi_+h_{\mu\nu}h^{\mu\nu}\rangle \\
& -67/2304\,n\,\langle\chi_+\rangle\langle h_{\mu\nu}h^{\mu\nu}\rangle + 3/128\,\langle A\chi_+^2\rangle - 3/64\,n^{-1}\,\langle A\chi_+\rangle\langle\chi_+\rangle \\
& -(3/128\,n^{-1}+1/128\,n)\langle A\rangle\langle\chi_+^2\rangle - (1/256 - 1/32\,n^{-2})\langle A\rangle\langle\chi_+\rangle^2 \\
& -(3/64-1/256\,n^2)\langle\chi_+u_\mu\chi_+u^\mu\rangle + (5/64\,n^{-1} - 1/256\,n)\langle\chi_+u_\mu\rangle^2 + 1/8\,n^{-2}\,\langle\chi_+^3\rangle \\
& -(1/8\,n^{-3}+1/16\,n^{-1})\langle\chi_+^2\rangle\langle\chi_+\rangle + (1/32\,n^{-4}+1/32\,n^{-2})\langle\chi_+\rangle^3 \\
& -(1/192-1/768\,n^2)i\langle\chi_-\{h_{\mu\nu},u^\mu u^\nu\}\rangle + 1/384\,n\,i\langle\chi_-h_{\mu\nu}\rangle\langle u^\mu u^\nu\rangle \\
& -(1/8\,n^{-1}+7/384\,n)i\langle h_{\mu\nu}u^\mu u^\nu\rangle\langle\chi_-\rangle + 13/96\,i\langle h_{\mu\nu}u^\mu\chi_-u^\nu\rangle \\
& +5/128\,n\,i\langle h_{\mu\nu}u^\mu\rangle\langle\chi_-u^\nu\rangle - (5/48-1/32\,n^{-2}-1/1152\,n^2)\langle A\chi_-^2\rangle \\
& +(3/32\,n^{-1}+19/1152\,n)\langle A\chi_-\rangle\langle\chi_-\rangle - (1/64\,n^{-1}+1/72\,n)\langle A\rangle\langle\chi_-^2\rangle \\
& +(31/2304-1/16\,n^{-2})\langle A\rangle\langle\chi_-\rangle^2 + (5/192-1/32\,n^{-2}-1/768\,n^2)\langle u_\mu\chi_-u^\mu\chi_-\rangle \\
& +(1/32\,n^{-1}-5/1152\,n)\langle u_\mu\chi_-\rangle^2 - (1/32-5/32\,n^{-2}+5/1152\,n^2)\langle\chi_-^2\chi_+\rangle \\
& -(1/16\,n^{-3}+29/1152\,n)\langle\chi_+\rangle\langle\chi_-^2\rangle - (1/16\,n^{-3}+1/16\,n^{-1}-7/288\,n)\langle\chi_+\chi_-\rangle\langle\chi_-\rangle \\
& +(1/48+1/32\,n^{-4}+1/64\,n^{-2})\langle\chi_+\rangle\langle\chi_-\rangle^2 + (1/64+1/192\,n^2)i\langle\chi_-\{\nabla_\mu\chi_+,u^\mu\}\rangle \\
& +(1/32\,n^{-1}-5/192\,n)i\langle\chi_-\rangle\langle\nabla_\mu\chi_+u^\mu\rangle - (1/32\,n^{-1}-11/192\,n)i\langle\nabla_\mu\chi_+\rangle\langle\chi_-u^\mu\rangle \\
& -(1/32-1/128\,n^2)\langle\nabla_\mu\chi_+\nabla^\mu\chi_+\rangle + 3/128\,n\,\langle\nabla_\mu\chi_+\rangle^2 \\
& +1/144\,n^2\,i\langle f_{+\mu\nu}\{\chi_+,u^\mu u^\nu\}\rangle + 1/9\,n\,i\langle\chi_+\rangle\langle f_{+\mu\nu}u^\mu u^\nu\rangle + 1/72\,n^2\,i\langle f_{+\mu\nu}u^\mu\chi_+u^\nu\rangle \\
& -(1/48-5/1152\,n^2)\langle f_{-\mu\nu}\,(h^{\nu\rho}u_\rho u^\mu + u^\mu u_\rho h^{\nu\rho})\rangle - 1/576\,n\,\langle f_{-\mu\nu}h^{\nu\rho}\rangle\langle u^\mu u_\rho\rangle \\
& +17/288\,n\,\langle f_{-\mu\nu}u^\mu\rangle\langle h^{\nu\rho}u_\rho\rangle + (1/48+1/192\,n^2)\langle f_{-\mu\nu}\,(u^\mu h^{\nu\rho}u_\rho + u_\rho h^{\nu\rho}u^\mu)\rangle \\
& -(1/16+1/2304\,n^2)i\langle f_{-\mu\nu}[\chi_-,u^\mu u^\nu]\rangle - 35/1152\,n\,i\langle f_{-\mu\nu}u^\nu\rangle\langle u^\mu\chi_-\rangle \\
& -5/288\,n^2\,\langle f_{-\mu\nu}\{\nabla^\mu\chi_+,u^\nu\}\rangle - 31/288\,n\,\langle\nabla^\mu\chi_+\rangle\langle f_{-\mu\nu}u^\nu\rangle \\
& -(1/96+1/288\,n^2)i\langle\nabla_\rho f_{+\mu\nu}[h^{\mu\rho},u^\nu]\rangle - (1/96+1/384\,n^2)i\langle\nabla^\mu f_{+\mu\nu}[h^{\nu\rho},u_\rho]\rangle\}
\end{aligned} \tag{8}$$

with $A = u_\mu u^\mu$ and $h_{\mu\nu} = \nabla_\mu u_\nu + \nabla_\nu u_\mu$. F is the meson decay constant in the chiral limit and the quantities u_μ, ∇_μ, $\chi_\pm$, $f_\pm^{\mu\nu}$ are defined as usual (e.g., in Ref. [7]).

In the chosen basis for the $SU(n)$ Lagrangian of $O(p^6)$, 109 of altogether 115 coefficients are divergent. For $n = 2$ or 3 [8], there are of course fewer independent terms but we postpone the explicit discussion of $\mathcal{L}_6$ for $n = 2, 3$ to a separate publication [3].

3. In this section, we sketch the main steps for extracting the double logs from the generating functional of $O(p^6)$. All technical details will be deferred to a separate publication [3].

In its most general form, the effective Lagrangian of $O(p^4)$ for chiral $SU(n)$ is given by

$$\begin{aligned}
\mathcal{L}_4 \;=\;& C_0\langle u_\mu u_\nu u^\mu u^\nu\rangle + C_1\langle u_\mu u^\mu\rangle^2 + C_2\langle u_\mu u_\nu\rangle\langle u^\mu u^\nu\rangle + C_3\langle u_\mu u^\mu u_\nu u^\nu\rangle \\
& + C_4\langle u_\mu u^\mu\rangle\langle\chi_+\rangle + C_5\langle u_\mu u^\mu\chi_+\rangle + C_6\langle\chi_+\rangle^2 + C_7\langle\chi_-\rangle^2 + \frac{1}{2}C_8\langle\chi_+^2 + \chi_-^2\rangle \\
& - iC_9\langle f_+^{\mu\nu}u_\mu u_\nu\rangle + \frac{1}{4}C_{10}\langle f_+^{\mu\nu}f_{+\mu\nu} - f_-^{\mu\nu}f_{-\mu\nu}\rangle \\
& + iC_{11}\langle\hat\chi_-(\nabla^\mu u_\mu - \frac{i}{2}\hat\chi_-)\rangle + C_{12}\langle(\nabla^\mu u_\mu - \frac{i}{2}\hat\chi_-)^2\rangle + \text{ contact terms} \qquad (9)
\end{aligned}$$

with $\hat\chi_- := \chi_- - \langle\chi_-\rangle/n$. The terms with coefficients C_{11}, C_{12} vanish at the classical solution defined by the equation of motion (EOM) $\nabla^\mu u_\mu - i\hat\chi_-/2 = 0$. All chiral Lagrangians of $O(p^4)$ can be written in the form (9) but there is of course some redundancy for $n = 2$ or 3. The Laurent expansion analogous to (3) is

$$C_i(d) = \mu^{d-4}\left[\frac{\Sigma_i}{(4\pi)^2(d-4)} + C_i^r(\mu) + \ldots\right] \qquad (10)$$

with known coefficients Σ_i [5].

The calculation of both reducible and one-loop diagrams in Fig. 1 proceeds along standard lines by expanding the chiral actions of $O(p^2)$ and $O(p^4)$ around the classical solution defined by the equation of motion. In particular, we need the expansion of S_4 up to second order in the fluctuation variables ξ defined in the usual way [5]:

$$S_4[\phi] = S_4[\phi_{\rm cl}] + S_{4i}[\phi_{\rm cl}]\xi_i + \frac{1}{2}S_{4ij}[\phi_{\rm cl}]\xi_i\xi_j + O(\xi^3) \qquad (11)$$

$$u(\phi) = u(\phi_{\rm cl})e^{i\xi(\phi)/2}, \quad \xi^\dagger = \xi, \quad \langle\xi\rangle = 0, \quad \xi(\phi_{\rm cl}) = 0, \quad \xi = \frac{1}{\sqrt{2}}\lambda_i\xi_i\,, \quad \langle\lambda_i\lambda_j\rangle = 2\delta_{ij}\,,$$

with λ_i the generators of $SU(n)$ in the fundamental representation.

We will not write down the explicit expressions for S_{4i} and S_{4ij} here (see Ref. [3]), but concentrate on the EOM terms in (9). As in the actual calculations, we switch to Euclidean space for the rest of this section. For investigating the influence of EOM terms, one needs the expansion

$$\nabla_\mu u_\mu + \frac{i}{2}\hat\chi_- = \frac{1}{\sqrt{2}}\lambda_i(-d_\mu d_\mu + \sigma)_{ij}\xi_j + O(\xi^2)\ . \qquad (12)$$

In (12), $(-d_\mu d_\mu + \sigma)_{ij} = G_{ij}^{-1}$ is the inverse of the full propagator[2] of $O(p^2)$ in the presence of external fields.

Turning first to the last term in (9), we infer from (12) that C_{12} cannot contribute to the reducible diagrams e,g in Fig. 1. Moreover, the one-loop diagram d proportional to C_{12} vanishes in dimensional regularization because of $\delta^d(0) = 0$. In other schemes, the one-loop contribution would be a quartic (and sixth-order) divergence that can be absorbed in $\mathcal{L}_6$. Chiral logs do not occur and we can set $C_{12} = 0$ without loss of generality.

---

[2] This is the propagator occurring in the functional diagrams of Fig. 1.

The situation is more subtle for C_{11}. The explicit factor G_{ij}^{-1} in (12) implies that all tree-level contributions from diagram g involving C_{11} are local, i.e. these terms can always be absorbed in $\mathcal{L}_6$. An explicit calculation shows in addition that diagrams d and e cancel exactly for the vertex associated with C_{11}. The final result is that C_{11} appears only in a (local) tree-level functional that has no bearing on chiral logs. Since we are free to choose any value for C_{11}, the choice $C_{11} = 0$ is the most convenient one for the calculation of the generating functional (for $n = 2, 3$, this corresponds to the Lagrangians of Refs. [9] and [5], respectively).

Diagrams c,e,g make up the reducible part of the generating functional of $O(p^6)$:

$$Z_6^{\mathrm{red}} = -\frac{1}{F^2}\left(S_{4i} + \frac{1}{2}D_{ikl}^{(3)}G_{kl}\right)G_{ij}\left(S_{4j} + \frac{1}{2}D_{jmn}^{(3)}G_{mn}\right) \tag{13}$$

where $D_{ikl}^{(3)}$ represents the cubic functional vertices in diagrams c,e (and b, for that matter). For $C_{11} = 0$, the combination

$$S_{4i} + \frac{1}{2}D_{ikl}^{(3)}G_{kl}$$

turns out to be finite and scale independent by itself (see also Ref. [10]). In other words, the one-loop divergences in c and e are cancelled by the divergent parts of the coefficients C_i $(i = 0, \ldots, 10)$ given in (10). This is precisely what one expects: the one-loop divergences are already taken care of by the renormalization at $O(p^4)$. Note however that this would not be the case for $C_{11} \neq 0$ in general.

Due to the finiteness and μ-independence of the reducible functional (13), the renormalized low-energy constants $C_i^r(\mu)$ always appear in the scale independent combination

$$C_i^r(\mu) - \frac{1}{2}\Sigma_i L \tag{14}$$

with L defined in (7). Therefore, the dependence of Z_6^{red} on chiral logs is completely fixed by the dependence on the $C_i^r(\mu)$ via diagram g. In the actual applications, chiral logs from this source will appear in mass and wave function renormalization, F_π, etc.[3]

Finally, the one-loop functional associated with diagram d is given by

$$Z_6^{L=1} = \frac{1}{F^2}S_{4ij}G_{ij}\ . \tag{15}$$

Decomposing the propagator $G_{ij}(x, y)$ in the usual way [11] into two parts, being singular and finite in the coincidence limit $x \to y$, respectively, $Z_6^{L=1}$ exhibits two types of divergences as sketched in Eq. (4). The nonlocal ones cancel with corresponding divergences from the irreducible diagrams a,b. The local divergences (of a,b and d) are cancelled by counterterms in $\mathcal{L}_6$. Following the arguments of the previous section, they also determine the finite parts given in (6). Keeping only the double-log contributions, the final result is expressed in terms of the double-log Lagrangian (8).

---

[3] In general, there can also be nonlocal double-log contributions. For instance, there will be such contributions for processes with six external mesons.

4. Let us now turn to some applications of the above results. For the case of two flavours, $n = 2$, a number of full two-loop calculations already exists (a short review can be found in [12]). We have checked that the double logarithms agree with those calculated for the following quantities: $\pi\pi$-scattering [2], F_π and M_π^2 [2, 13], radiative pion decay [14] and the pion vector and scalar form factors [15].

In the case of three flavours we agree with the double logarithms as calculated for the vector two-point function [16], for M_π^2, M_η^2, F_π and F_η [17]. To emphasize that these agreements provide nontrivial checks on our calculations, we show here the p^6 contributions from the k_i defined in (7) and from terms of the type $L_i^r L_j^r$ for F_π/F:

$$\begin{aligned} F_\pi^4(F_\pi/F\,)^{(6)} \;=\; & M_\pi^4(56L_4^{r2} + 112L_4^rL_5^r - 64L_4^rL_6^r - 64L_4^rL_8^r + 56L_5^{r2} - 64L_5^rL_6^r \\ & -64L_5^rL_8^r + 65/9\,k_1 + 73/18\,k_2 + 32/9\,k_3 - 20/3\,k_4 - 5\,k_5) + M_\pi^2M_K^2(224L_4^{r2} \\ & +160L_4^rL_5^r - 256L_4^rL_6^r - 128L_5^rL_6^r - 16/9\,k_1 - 4/9\,k_2 - 4/9\,k_3 - 37/6\,k_4 - 5/2\,k_5) \\ & +M_K^4(224L_4^{r2} + 64L_4^rL_5^r - 256L_4^rL_6^r - 128L_4^rL_8^r + 104/9\,k_1 + 26/9\,k_2 \\ & +61/18\,k_3 - 29/3\,k_4)\ . \end{aligned} \tag{16}$$

Here and in the following, as indicated by the superscript (6), we only display the partial results of $O(p^6)$ (in the sense explained above) for all quantities considered. In addition we always use isospin symmetry.

The result for F_K/F_π is new:

$$\begin{aligned} F_\pi^4(F_K/F_\pi)^{(6)} \;=\; & M_\pi^2(M_K^2 - M_\pi^2)(32L_4^rL_5^r + 40L_5^{r2} - 64L_5^rL_6^r - 64L_5^rL_8^r + k_1 \\ & +5/2\,k_2 + 19/12\,k_3 - 37/12\,k_5 + 4\,k_7 + 2\,k_8) + M_K^2(M_K^2 - M_\pi^2)(64L_4^rL_5^r + 24L_5^{r2} \\ & -128L_5^rL_6^r - 64L_5^rL_8^r + k_1 + 5/2\,k_2 + 23/12\,k_3 - 17/12\,k_5 - 4\,k_7 - 2\,k_8)\ . \end{aligned} \tag{17}$$

We defer a numerical discussion to the next section. The p^4 results for F_π/F and F_K/F_π can be found in [5].

We have also worked out the partial p^6 results for several relevant semileptonic kaon decays. For the two K_{l3} form factors f_+ and f_0 we obtain:

$$\begin{aligned} F_\pi^4(f_+(t))^{(6)} \;=\; & (M_K^2 - M_\pi^2)^2(8L_5^{r2} + 1/3\,k_1 + 1/6\,k_2 + 1/36\,k_3 - \ k_4) \\ & +tM_\pi^2(8L_5^rL_9^r - \ k_1 + 1/2\,k_2 - 1/4\,k_3 - 1/3\,k_4 - 1/2\,k_5 - \ k_9) \\ & +tM_K^2(-8L_5^rL_9^r - \ k_1 + 1/2\,k_2 - 5/4\,k_3 - 1/3\,k_4 - 1/2\,k_9) \\ & +t^2(1/3\,k_1 - 1/6\,k_2 + 1/4\,k_3 - 1/8\,k_9)\ , \end{aligned} \tag{18}$$

$$\begin{aligned} F_\pi^4(f_0(t))^{(6)} \;=\; & (M_K^2 - M_\pi^2)^2(8L_5^{r2} + 1/3\,k_1 + 1/6\,k_2 + 1/36\,k_3 - \ k_4) \\ & +tM_\pi^2(32L_4^rL_5^r - 64L_5^rL_6^r - 64L_5^rL_8^r + 48L_5^{r2} + 4/3\,k_1 + 11/3\,k_2 \\ & +22/9\,k_3 + \ k_4 - 23/6\,k_5 + 4\,k_7 + 2\,k_8) \\ & +tM_K^2(64L_4^rL_5^r - 128L_5^rL_6^r - 64L_5^rL_8^r + 16L_5^{r2} + 4/3\,k_1 + 11/3\,k_2 \\ & +25/9\,k_3 + \ k_4 - 2/3\,k_5 - 4\,k_7 - 2\,k_8) \\ & +t^2(-2/3\,k_1 - 4/3\,k_2 - 8/9\,k_3 - 3/4\,k_5)\ . \end{aligned} \tag{19}$$

The Ademollo-Gatto theorem [18] is the reason for the appearance of the factor $(M_K^2 - M_\pi^2)^2$. The definition of the two form factors and a discussion of experimental results can be found in [19]. The p^4 results can also be found there and were first obtained in [20].

We also derived the corresponding results for the K_{e4} form factors F and G. The definition of all quantities appearing here can be found in [19, 21] and the p^4 results using the same notation can also be found there. They were first derived in [22, 23].

$$
\begin{aligned}
(\sqrt{2}F_\pi/M_K)F_\pi^4\, F(q^2,\nu,s_l)^{(6)} \;=\;& M_\pi^4 q^4(-404\,k_1 - 178\,k_2 - 431/3\,k_3) \\
&+M_\pi^4 q^2(512L_1^rL_5^r + 128L_2^rL_5^r + 160L_3^rL_5^r - 1904/3\,k_1 - 2686/9\,k_2 - 11987/54\,k_3 \\
&-80\,k_4 - 27\,k_5) + M_\pi^4(256L_1^rL_5^r + 128L_2^rL_5^r + 96L_3^rL_5^r + 256L_4^{r2} + 416L_4^rL_5^r \\
&-512L_4^rL_6^r - 512L_4^rL_8^r + 56L_5^{r2} - 64L_5^rL_6^r - 64L_5^rL_8^r - 1969/9\,k_1 \\
&-2509/18\,k_2 - 9133/108\,k_3 - 854/9\,k_4 - 271/12\,k_5 - 16\,k_6 - 20\,k_7 - 18\,k_8) \\
&+M_\pi^2M_K^2q^2(-512L_1^rL_5^r - 128L_2^rL_5^r - 160L_3^rL_5^r + 437/3\,k_1 + 229/6\,k_2 \\
&+691/12\,k_3 - 52\,k_4) + M_\pi^2M_K^2(-256L_1^rL_5^r - 96L_2^rL_5^r - 88L_3^rL_5^r + 512L_4^{r2} \\
&-64L_4^rL_5^r - 1024L_4^rL_6^r - 32L_5^{r2} - 128L_5^rL_6^r + 1000/9\,k_1 + 247/18\,k_2 \\
&+9005/216\,k_3 - 100/9\,k_4 + 73/12\,k_5 - 44\,k_6 + 20\,k_7 - 12\,k_8) \\
&+M_\pi^2 s_l q^2(\,k_1 + 71/6\,k_2 - 5/12\,k_3 - 9/2\,k_9) + M_\pi^2 s_l(-32L_2^rL_5^r - 8L_3^rL_5^r \\
&+8L_5^rL_9^r + 4/3\,k_1 + 337/18\,k_2 + 319/216\,k_3 + 3/2\,k_5 - 43/8\,k_9) \\
&+M_\pi^2\nu q^2(-5\,k_1 - 15/2\,k_2 - 19/4\,k_3) + M_\pi^2\nu(-8L_3^rL_5^r - 9/2\,k_1 - 33/4\,k_2 \\
&-19/6\,k_3 - 2\,k_4) + M_K^4(-32L_2^rL_5^r - 8L_3^rL_5^r + 8L_5^{r2} - 1/3\,k_1 - 6\,k_2 - 61/72\,k_3 \\
&-4/3\,k_4) + M_K^2 s_l(32L_2^rL_5^r + 8L_3^rL_5^r - 8L_5^rL_9^r + 41/6\,k_2 - 11/24\,k_3 - 7/8\,k_9) \\
&+M_K^2\nu(8L_3^rL_5^r - 19/6\,k_1 - 17/6\,k_2 - 145/72\,k_3 - \;k_4) + s_l^2(-1/3\,k_2 + 1/3\,k_3 \\
&+1/4\,k_9) + s_l\nu(\,k_1 + 7/4\,k_2 + 9/8\,k_3) + \nu^2(-5/12\,k_2 + 7/24\,k_3)\;,
\end{aligned}
\tag{20}
$$

$$
\begin{aligned}
(\sqrt{2}F_\pi/M_K)F_\pi^4\, G(q^2,\nu,s_l)^{(6)} \;=\;& M_\pi^4 q^4(-20\,k_2 - 2\,k_3) \\
&+M_\pi^4 q^2(-32L_3^rL_5^r - 22/3\,k_1 - 119/3\,k_2 - 34/9\,k_3 - 8\,k_4) + M_\pi^4(-32L_3^rL_5^r \\
&+32L_4^rL_5^r + 56L_5^{r2} - 64L_5^rL_6^r - 64L_5^rL_8^r - 19/3\,k_1 - 103/6\,k_2 - 7/36\,k_3 - 8\,k_4 \\
&-55/12\,k_5 + 4\,k_7 + 2\,k_8) + M_\pi^2M_K^2q^2(32L_3^rL_5^r - 25/3\,k_1 - 73/6\,k_2 - 205/36\,k_3 \\
&-4\,k_4) + M_\pi^2M_K^2(24L_3^rL_5^r + 64L_4^rL_5^r - 32L_5^{r2} - 128L_5^rL_6^r - 23/2\,k_1 - 169/12\,k_2 \\
&-181/36\,k_3 + 2\,k_4 + 7/3\,k_5 - 12\,k_6 - 4\,k_7 - 8\,k_8) + M_\pi^2 s_l q^2(7/3\,k_1 + 47/6\,k_2 \\
&+13/4\,k_3) + M_\pi^2 s_l(8L_3^rL_5^r + 8L_5^rL_9^r + 13/6\,k_1 + 35/4\,k_2 + 79/36\,k_3 + 4/3\,k_4 \\
&-3/4\,k_5 - 11/8\,k_9) + M_\pi^2\nu q^2(-5/3\,k_1 - 5/2\,k_2 + 13/12\,k_3) + M_\pi^2\nu(32L_2^rL_5^r \\
&+8L_3^rL_5^r - 7/3\,k_1 - 10\,k_2 - 35/72\,k_3 - 2/3\,k_4 - 9/4\,k_5) \\
&+M_K^4(8L_3^rL_5^r + 8L_5^{r2} - 29/12\,k_1 - 13/8\,k_2 - 223/144\,k_3 - 2\,k_4) + M_K^2 s_l(-8L_3^rL_5^r \\
&-8L_5^rL_9^r + 7/3\,k_1 + 3\,k_2 + 73/72\,k_3 + 2/3\,k_4 - 1/2\,k_9) + M_K^2\nu(-32L_2^rL_5^r - 8L_3^rL_5^r \\
&-2/3\,k_1 - 79/12\,k_2 - 7/12\,k_3 - 1/3\,k_4) + s_l^2(-1/4\,k_1 - 7/8\,k_2 - 7/16\,k_3 - 1/8\,k_9) \\
&+s_l\nu(1/3\,k_1 + 7/12\,k_2 - 3/8\,k_3 - 3/8\,k_9) + \nu^2(-5/12\,k_1 - 25/24\,k_2 - 7/16\,k_3)\;.
\end{aligned}
\tag{21}
$$

change	set A	$\mu = 0.9$ GeV	set B	$M = 0.26$ GeV
$(F_\pi/F)^{(6)}$	0.013	0.029	-0.050	-0.005
$(F_K/F_\pi)^{(6)}$	0.08	0.12	0.06	0.24
$10^3 \cdot f_{+1}$	3.6	-4.8	2.0	-2.4
$f_{+2}[\mathrm{GeV}^{-2}]$	-0.28	-0.24	-0.25	-0.26
$f_{+3}[\mathrm{GeV}^{-4}]$	0.58	0.77	0.55	1.5
$f_{02}[\mathrm{GeV}^{-2}]$	0.30	0.51	0.18	0.74
$f_{03}[\mathrm{GeV}^{-4}]$	0.26	0.23	0.41	1.54
F_1	0.86	1.13	0.75	2.5
F_2	0.38	0.51	0.17	0.74
F_3	0.000	-0.019	0.052	0.14
G_1	-0.15	-0.20	-0.04	0.18
G_2	0.006	0.014	0.035	0.17
G_3	0.010	0.012	0.011	0.031

Table 1: Numerical results for the partial p^6 corrections for several quantities. See text for definitions and input parameters.

5. We will now show some numerical results using the previous formulas. It should be kept in mind that the final numbers are quite sensitive to the input numbers and that there are several uncertainties:

1. The logarithms in the k_i can have varying scales in them, i.e. the scale M in Eq. (7) can be varied. Since all the numerical examples refer to chiral $SU(3)$, the choice $M = M_K$ is the most natural one. We will contrast the results for $M = M_K$ with those for $M = \sqrt{M_K M_\pi}$.
2. The final results are of course μ dependent. Both the values of $L_i^r(\mu)$ and the value of L change as functions of μ.
3. The values of $L_i^r(\mu)$ that are used as input are in general only determined via p^4 calculations.
4. The parts we included are dominant when L is large. For the $n = 3$ calculations considered here, the logarithm typically is not as dominant as for $n = 2$. Thus the remaining parts of the loop amplitudes can be important.
5. The contributions from the p^6 Lagrangian are all set to zero.

For all these reasons the numbers quoted here should only be used as an indication of the size of the p^6 corrections. As input values we have used $F_\pi =$

92.4 MeV, $M_K = 495$ MeV, $M_\pi = 135$ MeV and $10^3 \cdot L^r_{(4,5,6,8,9)}(0.77 \text{ GeV}) = (-0.3, 1.4, -0.2, 0.9, 6.9)$. We will vary μ, the scale M and use two sets of $L^r_{(1,2,3)}$. The first set corresponds to the unitarized fit of [21], $10^3 \cdot L^r_{(1,2,3)}(0.77 \text{ GeV}) = (0.4, 1.35, -3.5)$ (set A) whereas $10^3 \cdot L^r_{(1,2,3)}(0.77 \text{ GeV}) = (0.6, 1.5, -3.3)$ (set B) is from the one-loop fit of the same reference. The values of $\mu = 0.77$ GeV, $M = M_K = 0.495$ GeV and set A are used unless otherwise shown. We write for the K_{l3} form factors

$$(f_+(t))^{(6)} = f_{+1} + f_{+2}t + f_{+3}t^2, \quad (f_0(t))^{(6)} = f_{01} + f_{02}t + f_{03}t^2 \tag{22}$$

and we evaluate the K_{e4} form factors at $s_l = \nu = 0$:

$$F^{(6)}(q^2, 0, 0) = F_1 + F_2 q^2 + F_3 q^4, \quad G^{(6)}(q^2, 0, 0) = G_1 + G_2 q^2 + G_3 q^4 \tag{23}$$

The results are shown in Table 1. Since by definition of the K_{l3} form factors $f_{01} = f_{+1}$ we do not display f_{01}. The last column shows that $M = \sqrt{M_K M_\pi} = 260$ MeV is probably not a good choice for terms of the type $\left(M_K^2 \ln(M/\mu)\right)^2$ leading to unreasonably big numbers in most cases. Not surprisingly, the numbers become even more unreasonable for $M = M_\pi$.

The general size of the correction to F_π/F is similar to the full result for the two-flavour case obtained in [15]. The correction to F_K/F_π is rather large and will, if the full p^6 calculation is of similar size, require a revision of the value for L^r_5.

As for the K_{l3} form factors, the small correction f_{+1} to $f_+(0)$ is an important result. The form factor at $t = 0$ is used as input in the determination of the Cabibbo angle from K_{l3} decays. It is therefore important to know that $f_+(0)$ is well protected from p^6 corrections [24]. The corrections to the slopes λ_+, λ_0 (f_{+2}, f_{02} in the table) have about the same size but opposite signs. These corrections are within the expectations for chiral $SU(3)$. However, the scalar slope might be more sensitive to p^6 corrections because the p^4 prediction for λ_0 is smaller than for λ_+ [20]. Finally, the curvatures induced at $O(p^6)$ are completely negligible in the physical region, in agreement with the observed Dalitz plot distributions and with theoretical expectations [20].

The size of the corrections to the K_{e4} form factors is such that a full p^6 calculation is desirable in order to refit L_1, L_2 and L_3. This is especially true for the slope of the form factor F (F_2 in the table) where the p^6 correction is of the same size as the observed slope [25]. Using the other inputs as above and fitting $L_{(1,2,3)}$ to $F(0,0,0) = 5.59 \pm 0.14$, $G(0,0,0) = 4.77 \pm 0.27$ and $(F(q^2,0,0) - F(0,0,0))/q^2|_{q^2=0.1} = (0.08 \pm 0.02)\, F(0,0,0)$ leads to $10^3 \cdot L^r_{(1,2,3)} = (0.25 \pm 0.3, 1.24 \pm 0.3, -4.75 \pm 1.9)$ to be compared with the one-loop K_{e4}-only fit of [21] with the same inputs of $(0.65 \pm 0.3, 1.63 \pm 0.28, -3.4 \pm 1.0)$.

6. We have determined the full double-pole divergence structure of chiral perturbation theory at $O(p^6)$. We have described the general method for extracting all contributions with double chiral logs (L^2) as well as the full dependence on $L^r_i \times L$ and $L^r_i \times L^r_j$. These partial p^6 corrections were then calculated for several quantities of physical interest. The corrections to F_K/F_π and to the slopes of the K_{l3} form factors are of the size expected for chiral $SU(3)$. On the other hand, we find small corrections to the K_{l3} form factor f_+ at $t = 0$, important for the determination of V_{us}. The corrections for K_{e4} decays are significant suggesting possible shifts in the values of $L^r_{(1,2,3)}$.

Acknowledgements We thank J. Gasser for participation in the early stages of this work and for continuous encouragement.

References

[1] G. Colangelo, Phys. Lett. B 350 (1995) 85; Phys. Lett. B 361 (1995) 234 (E).

[2] J. Bijnens, G. Colangelo, G. Ecker, J. Gasser and M.E. Sainio, Phys. Lett. B 374 (1996) 210; Nucl. Phys. B 508 (1997) 263.

[3] J. Bijnens, G. Colangelo and G. Ecker, in preparation.

[4] E.g., J.C. Collins, Renormalization (Cambridge University Press, Cambridge, 1984).

[5] J. Gasser and H. Leutwyler, Nucl. Phys. B 250 (1985) 465.

[6] S. Weinberg, Physica 96A (1979) 327.

[7] G. Ecker, J. Gasser, A. Pich and E. de Rafael, Nucl. Phys. B 321 (1989) 311.

[8] H.W. Fearing and S. Scherer, Phys. Rev. D 53 (1996) 315.

[9] J. Gasser, M.E. Sainio and A. Švarc, Nucl. Phys. B 307 (1988) 779.

[10] G. Ecker and M. Mojžiš, Phys. Lett. B 365 (1996) 312.

[11] I. Jack and H. Osborn, Nucl. Phys. B 207 (1982) 474.

[12] J. Bijnens, hep-ph/9710341.

[13] U. Bürgi, Phys. Lett. B 377 (1996) 147; Nucl. Phys. B 479 (1996) 392.

[14] J. Bijnens and P. Talavera, Nucl. Phys. B 489 (1997) 387.

[15] J. Bijnens, G. Colangelo and P. Talavera, JHEP 05 (1998) 014

[16] E. Golowich and J. Kambor, Nucl. Phys. B 447 (1995) 373.

[17] E. Golowich and J. Kambor, Phys. Rev. D 58 (1998) 3604.

[18] M. Ademollo and R. Gatto, Phys. Rev. Lett. 13 (1964) 264.

[19] J. Bijnens et al., hep-ph/9411311, chapter 7.1 in “The Second DAΦNE Physics Handbook”, eds. L. Maiani, G. Pancheri and N. Paver (INFN, Frascati, 1995).

[20] J. Gasser and H. Leutwyler, Nucl. Phys. B 250 (1985) 517.

[21] J. Bijnens, G. Colangelo and J. Gasser, Nucl. Phys. B 427 (1994) 427.

[22] J. Bijnens, Nucl. Phys. B 337 (1990) 635.

[23] C. Riggenbach et al., Phys. Rev. D 43 (1991) 127.

[24] H. Leutwyler and M. Roos, Z. Phys. C 25 (1984) 91.

[25] L. Rosselet et al., Phys. Rev. D 15 (1977) 574.